

**Title: Fructosamine; is the current interest in alternative glycaemic markers justified?**

Running head: A role for fructosamine?

KE Shipman<sup>1</sup>, M Jawad<sup>1</sup>, KM Sullivan<sup>1</sup>, C Ford<sup>1</sup>, R Gama<sup>1,2</sup>

<sup>1</sup>New Cross Hospital, Clinical Chemistry, Wolverhampton WV10 0QP, UK.

Phone: 01902 695290

Fax: 01902 695618

<sup>2</sup>Research Institute, Healthcare Sciences, Wolverhampton University, Wolverhampton, UK.

Corresponding Author: Dr Kate Shipman, *[kate.shipman@doctors.net.uk](mailto:kate.shipman@doctors.net.uk)*

Word count: 329

Funding sources: No sources of funding to declare.

Conflicts of interest: The authors of this letter have no conflicts of interest to disclose.

Sir

**Fructosamine; is the current interest in alternative glycaemic markers justified?**

The interest in alternative glycaemic markers, such as fructosamine, has been rekindled due to increasing data on their ability to improve diagnosis [1] and predict onset and complications [2] of diabetes. Further clinical validation of these markers is called for [3] however the limitations must also be elucidated so, rather like HbA1c, the non-glycaemic physiological, pathological and analytical variables become well known.

Many of the variables that affect HbA1c appear to similarly affect fructosamine despite the different physiological compartments. The most commonly known non-glycaemic factor affecting fructosamine is albumin [3] although data on this is conflicting. We recently reported a multivariable univariate linear regression model of fructosamine (Table 3) [4] in which the effect of ethnicity (South Asian versus Caucasian), sex, age, presence of chronic kidney disease (CKD), vitamin B12, ferritin, folate, haemoglobin, mean cell volume (MCV), fasting glucose, HbA1c, albumin and C reactive protein (CRP) were examined [4]. The significant affect of albumin levels was confirmed as were the effects of ethnicity, age, presence of CKD (primarily CKD stage 3 versus CKD <3), vitamin B12, folate, albumin, CRP, fasting glucose levels, HbA1c and haemoglobin. Ferritin, MCV and sex were the only non-significant variables [4].

The large number of significant variables had not been anticipated, included due to their established effect on HbA1c, confirming our current poor grasp of fructosamine metabolism. Body mass index is also negatively associated with fructosamine in those without diabetes [5] and falls during pregnancy [6] further complicating its interpretation. Thyroid status may

affect fructosamine interpretation [3] however not necessarily independently of deranged glucose and protein metabolism [7].

We, therefore, read with interest the new data but would urge caution on the widespread adoption of fructosamine until the significance of non-glycaemic variables can be established in the clinical field. In particular HbA1c remains the biomarker of choice for monitoring glycaemic control in those with CKD [8], whereas fructosamine should be used with caution [4].

Yours sincerely

Kate E Shipman<sup>1</sup>, Mohammed Jawad<sup>1</sup>, Katie M Sullivan<sup>1</sup>, Clare Ford<sup>1</sup>, Rousseau Gama<sup>1,2</sup>

<sup>1</sup>Clinical Chemistry, New Cross Hospital, Wolverhampton, West Midlands and <sup>2</sup>Research Institute, Healthcare Sciences, Wolverhampton University, Wolverhampton, West Midlands, UK.

## REFERENCES

1. Malmström H, Walldius G, Grill V, Jungner I, Gudbjörnsdottir S, Hammar N. Fructosamine is a useful indicator of hyperglycaemia and glucose control in clinical and epidemiological studies--cross-sectional and longitudinal experience from the AMORIS cohort. *PLoS One* 2014 29;9(10):e111463.
2. Selvin E, Rawlings AM, Grams M, Klein R, Sharrett AR, Steffes M, Coresh J. Fructosamine and glycated albumin for risk stratification and prediction of incident diabetes and microvascular complications: a prospective cohort analysis of the Atherosclerosis Risk in Communities (ARIC) study. *Lancet Diabetes Endocrinol* 2014;2(4):279-88.
3. Parrinello CM, Selvin E. Beyond HbA1c and glucose: the role of nontraditional glycemic markers in diabetes diagnosis, prognosis, and management. *Curr Diab Rep* 2014;14(11):548.
4. Shipman KE, Jawad M, Sullivan KM, Ford C, Gama R. Ethnic/racial determinants of glycemic markers in a UK sample. *Acta Diabetol* 2015 [Epub ahead of print].
5. Poon AK, Juraschek SP, Ballantyne CM, Steffes MW, Selvin E. Comparative associations of diabetes risk factors with five measures of hyperglycemia. *BMJ Open Diabetes Res Care* 2014;2(1):e000002.

6. Hartland AJ, Smith JM, Clark PM, Webber J, Chowdhury T, Dunne F. Establishing trimester- and ethnic group-related reference ranges for fructosamine and HbA1c in non-diabetic pregnant women. *Ann Clin Biochem* 1999;36:235-7.
7. Udupa SV, Manjrekar PA, Udupa VA, Vivian D. Altered fructosamine and lipid fractions in subclinical hypothyroidism. *J Clin Diagn Res* 2013;7(1):18-22.
8. Shipman KE, Jawad M, Sullivan KM, Ford C, Gama R. Effect of Chronic Kidney Disease on A1C in Individuals Being Screened for Diabetes. *Prim Care Diab* 2014;pii:S1751-9918(14)00053-9.